

AD-A063 978

WISCONSIN UNIV-MADISON MATHEMATICS RESEARCH CENTER

F/G 12/1

REMARKS ON THE SCHROEDINGER OPERATOR WITH SINGULAR COMPLEX POTE--ETC(U)

AUG 78 H BREZIS, T KATO

DAAG29-75-C-0024

UNCLASSIFIED

MRC-TSR-1875

NL

1 OF 1
ADA
063978



END
DATE
FILMED

3-79
DDC

AD A063978

MRC Technical Summary Report #1875

REMARKS ON THE SCHRÖDINGER OPERATOR
WITH SINGULAR COMPLEX POTENTIALS


Haïm Brezis and Tosio Kato

LEVEL II

Mathematics Research Center
University of Wisconsin-Madison
610 Walnut Street
Madison, Wisconsin 53706

August 1978

(Received June 7, 1978)



Approved for public release
Distribution unlimited

Sponsored by

U. S. Army Research Office
P. O. Box 12211
Research Triangle Park
North Carolina 27709

National Science Foundation
Washington, D. C. 20550

79 01 30 042

DDC FILE COPY

UNIVERSITY OF WISCONSIN - MADISON
MATHEMATICS RESEARCH CENTER

6 REMARKS ON THE SCHRÖDINGER OPERATOR WITH SINGULAR
COMPLEX POTENTIALS.

10 Haim/Brezis⁹³ and Tosio/Kato⁹⁴

9 Technical Summary Report, 1875
August 1978

11 Aug 78

ABSTRACT

12 22 p.

Let $A = -\Delta + V(x)$ be a Schrödinger operator on an (arbitrary) open set $\Omega \subset \mathbb{R}^m$, where $V \in L^1_{\text{loc}}(\Omega)$ is a complex valued function. We consider the "maximal" realization of A in $L^2(\Omega)$ under Dirichlet boundary condition, that is

$$D(A) = \{u \in H^1_0(\Omega); Vu \in L^1_{\text{loc}}(\Omega) \text{ and } -\Delta u + Vu \in L^2(\Omega)\}.$$

When $\Omega = \mathbb{R}^m$ we also consider the operator

$$A_1 = -\Delta + V$$

with domain

$$D(A_1) = \{u \in L^2(\Omega); Vu \in L^1_{\text{loc}}(\Omega) \text{ and } -\Delta u + Vu \in L^2(\Omega)\}.$$

A special case of our main results is:

Theorem: Let $m \geq 3$; assume that the function $\max\{-\operatorname{Re} V, 0\}$ belongs to $L^\infty(\Omega) + L^{m/2}(\Omega)$ and also to $L^{(m/2)+\epsilon}_{\text{loc}}(\Omega)$ for some $\epsilon > 0$. Then A (resp. A_1) is closable and $\bar{A} + \lambda$ (resp. $\bar{A}_1 + \lambda$) is m -accretive for some real constant λ .

AMS (MOS) Subject Classifications: 35J10, 47B44

Key Words: Schrödinger operator, Complex potentials, m -accretive operator

Work Unit Number 1 (Applied Analysis)

15 DAAG-29-75-C-0024 NSF-MCS 76-04655

page - i -

¹Dept. de Mathématiques, Université Paris VI, 4 pl. Jussieu, 75230 Paris 05, France.

²Dept. of Mathematics, University of California, Berkeley, CA 94720.

Sponsored by

3) the United States Army under Contract No. DAAG29-75-C-0024;

4) the National Science Foundation under Grant No. MCS76-04655.

224 200

alt

SIGNIFICANCE AND EXPLANATION

∇ ϵ
 Schrödinger operators of the form $A = -\Delta + V(x)$, where Δ is the Laplacian and V is a scalar potential, arise in quantum mechanics and other areas. Delicate questions concerning what domain should be assigned to A must be settled in order to have a good theory. These questions are answered here for a very general class of potentials V which may even have complex values.

ACCESSION for	
NTIS	White Section <input checked="" type="checkbox"/>
DDC	Buff Section <input type="checkbox"/>
UNANNOUNCED	<input type="checkbox"/>
JUSTIFICATION	
BY	
DISTRIBUTION/AVAILABILITY CODES	
Dist.	SPECIAL

The responsibilities for the wording and views expressed in this descriptive summary lies with MRC, and not with the authors of this report.

page - ii -

REMARKS ON THE SCHRÖDINGER OPERATOR WITH SINGULAR COMPLEX POTENTIALS

Haim Brezis^{1,3} and Tosio Kato^{2,4}

1. Introduction

Let $A = -\Delta + V(x)$ be a Schrödinger operator on an (arbitrary) open set $\Omega \subset \mathbb{R}^m$, where $V \in L^1_{\text{loc}}(\Omega)$ is a complex valued function. We consider the "maximal" realization of A in $L^2(\Omega)$ under Dirichlet boundary condition, that is

$$D(A) = \{u \in H^1_0(\Omega); Vu \in L^1_{\text{loc}}(\Omega) \text{ and } -\Delta u + Vu \in L^2(\Omega)\}.$$

When $\Omega = \mathbb{R}^m$ we also consider the operator

$$A_1 = -\Delta + V$$

with domain

$$D(A_1) = \{u \in L^2(\Omega); Vu \in L^1_{\text{loc}}(\Omega) \text{ and } -\Delta u + Vu \in L^2(\Omega)\}.$$

We state now our main results (see Theorems 3.1 and 3.2) in a special case.

Theorem: Let $m \geq 3$; assume that the function $\max\{-\operatorname{Re} V, 0\}$ belongs to $L^\infty(\Omega) + L^{m/2}(\Omega)$ and also to $L^{(m/2)+\varepsilon}_{\text{loc}}(\Omega)$ for some $\varepsilon > 0$. Then A (resp. A_1) is closable and $\bar{A} + \lambda$ (resp. $\bar{A}_1 + \lambda$) is m -accretive for some real constant λ .

We emphasize the fact that $\max\{\operatorname{Re} V, 0\}$ and $\operatorname{Im} V$ could be arbitrary functions in $L^1_{\text{loc}}(\Omega)$.

Our methods rely on some measure theoretic arguments and standard techniques of DeGiorgi-Moser-Stampacchia type, related to the weak form of the maximum principle.

The distributional inequality

$$\Delta|u| \geq \operatorname{Re}[\Delta u \operatorname{sign} \bar{u}]$$

proved in [3] plays a crucial role. We also use a result from [1] concerning a property of Sobolev spaces.

In order to describe our method in a simple case we begin in Section 2 with real valued potentials. The main results in Section 2 are essentially known (see [3], [4], [8]) - except perhaps for Theorem 2.2 when $m \leq 4$.

¹Dept. de Mathématiques, Université Paris VI, 4 pl. Jussieu, 75230 Paris 05, France.

²Dept. of Mathematics, University of California, Berkeley, CA 94720.

Sponsored by

3) the United States Army under Contract No. DAAG29-75-C-0024;

4) the National Science Foundation under Grant No. MCS76-04655.

In Section 3 we turn to the case of complex potentials. Schrödinger operators with complex potentials have been studied by Nelson [6]. His results were extended in [5]. Here we allow more general singularities.

We thank Professors R. Jensen and B. Simon for useful suggestions and discussions (with the first author).

2. Real valued potentials

Let Ω be an (arbitrary) open subset of \mathbb{R}^m and let $H = L^2 = L^2(\Omega; \mathbb{C})$. Let $q \in L^1_{\text{loc}}(\Omega)$ be a real valued function. Set

$$q^+ = \max(q, 0), \quad q^- = \max(-q, 0).$$

Assume

$$(1) \quad q^- \in L^\infty(\Omega) + L^p(\Omega)$$

with

$$\begin{cases} p = \frac{m}{2} & \text{when } m \geq 3 \\ p > 1 & \text{when } m = 2 \\ p = 1 & \text{when } m = 1. \end{cases}$$

Consider the operator A defined in H by

$$A = -\Delta + q(x)$$

with

$$D(A) = \{u \in H^1_0(\Omega); qu \in L^1_{\text{loc}}(\Omega) \text{ and } -\Delta u + qu \in L^2(\Omega)\}.$$

The main results are the following:

Theorem 2.1. A is self-adjoint and $A + \lambda_1$ is m -accretive for some real constant λ_1 .

Furthermore $u, v \in D(A)$ imply $q|u|^2 \in L^1(\Omega)$, $q|v|^2 \in L^1(\Omega)$ and

$$(2) \quad (Au, v) = \int \text{grad} u \cdot \text{grad} \bar{v} + \int q u \bar{v}.$$

When $\Omega = \mathbb{R}^m$ we also consider the operator A_1 defined in H by

$$A_1 = -\Delta + q(x)$$

with

$$D(A_1) = \{u \in L^2(\Omega); qu \in L^1_{\text{loc}}(\Omega) \text{ and } -\Delta u + qu \in L^2(\Omega)\}.$$

Only when $m = 3$ or $m = 4$ we will make the additional assumption:

$$(3) \quad q^- \in L^{p+\epsilon}_{\text{loc}}(\Omega) \text{ with } p = \frac{3}{2} \text{ when } m = 3 \text{ and } p = 2 \text{ when } m = 4, \text{ for some arbitrarily small } \epsilon > 0.$$

More precisely we assume that for each $x_0 \in \mathbb{R}^m$ there exists a neighborhood U of x_0 and some $\epsilon > 0$ (depending on x_0) such that $q^- \in L^{p+\epsilon}(U)$.

Theorem 2.2: Under the assumptions (1) and (3), $A_1 = A$.

Our first lemma is well known:

Lemma 2.1: Assume (1). Then for every $\varepsilon > 0$, there exists a constant λ_ε such that

$$\int q^- |u|^2 \leq \varepsilon \|\text{gradu}\|_{L^2}^2 + \lambda_\varepsilon \|u\|_{L^2}^2 \quad \forall u \in H_0^1(\Omega).$$

In particular

$$\int q^- |u|^2 \leq \|\text{gradu}\|_{L^2}^2 + \lambda_1 \|u\|_{L^2}^2 \quad \forall u \in H_0^1(\Omega).$$

Proof: Write $q^- = q_1 + q_2$ with $q_1 \in L^\infty(\Omega)$ and $q_2 \in L^p(\Omega)$. Then for each $k > 0$ we have

$$\begin{aligned} \int q^- |u|^2 &\leq \|q_1\|_{L^\infty} \|u\|_{L^2}^2 + \int_{\{|q_2| > k\}} |q_2| |u|^2 + k \int_{\{|q_2| \leq k\}} |u|^2 \\ &\leq (\|q_1\|_{L^\infty} + k) \|u\|_{L^2}^2 + \|q_2\|_{L^p(\{|q_2| > k\})} \|u\|_{L^t}^2 \end{aligned}$$

with

$$\frac{1}{p} + \frac{2}{t} = 1.$$

In case $m \geq 3$ we find $t = 2^*$ where 2^* is the Sobolev exponent, that is $\frac{1}{2^*} = \frac{1}{2} - \frac{1}{m}$.

By the Sobolev imbedding theorem we have

$$\|u\|_{L^t} \leq C \|\text{gradu}\|_{L^2} \quad \forall u \in H_0^1(\Omega).$$

When $m = 2$ we find $2 < t < \infty$ and it is known that

$$\|u\|_{L^t} \leq C(\|\text{gradu}\|_{L^2} + \|u\|_{L^2}) \quad \forall u \in H_0^1(\Omega).$$

When $m = 1$ we find $t = \infty$ and it is known that

$$\|u\|_{L^\infty} \leq C(\|\text{gradu}\|_{L^2} + \|u\|_{L^2}) \quad \forall u \in H_0^1(\Omega).$$

We reach the conclusion of Lemma 2.1 in all the cases by choosing k large enough so that

$$C^2 \|q_2\|_{L^p(\{|q_2| > k\})} < \varepsilon.$$

Remark 2.1: Assumption (1) is used in all the results of this paper only through

Lemma 2.1 and it may in fact be weakened to a "locally uniform L^p -condition":

$$(1') \quad \|q^-\|_{L^p(\Omega \cap B_r(y))} \rightarrow 0 \text{ as } r \rightarrow 0 \text{ uniformly in } y \in \Omega,$$

where

$$B_r(y) = \{x \in \mathbb{R}^m; |x - y| \leq r\}.$$

Indeed let $\varphi \in \mathcal{D}_+(\mathbb{R}^m)$ with $\text{supp } \varphi \subset B_r(0)$ and $\|\varphi\|_{L^2} = 1$. Then, writing $\varphi_y(x) = \varphi(x - y)$,

$$\int q^- |u|^2 = \int dy \int q^- |u \varphi_y|^2 \leq \int \|q^-\|_{L^p(B_r(y))} \|\varphi_y\|_{L^2}^2 dy.$$

Here $\|q^-\|_{L^p(B_r(y))} \leq \delta$ for any small δ by (1') if r is chosen small. So

$$\begin{aligned} \int q^- |u|^2 &\leq \delta \int \|\varphi_y\|_{L^2}^2 dy \leq C\delta \int \|\text{grad}(u \varphi_y)\|_{L^2}^2 dy \\ &\leq 2C\delta \int (\|\varphi_y \text{grad} u\|_{L^2}^2 + \|u \text{grad} \varphi_y\|_{L^2}^2) dy \\ &= 2C\delta (\|\text{grad} u\|_{L^2}^2 + C_r \|u\|_{L^2}^2). \end{aligned}$$

Choosing δ so that $2C\delta = \varepsilon$, one gets the conclusion of Lemma 2.1. Such a locally uniform L^p -condition was used by Simader [7].

We recall a result of [1] which will be used in the proof of Theorem 2.1⁽¹⁾.

Lemma 2.2: Let $T \in H^{-1}(\Omega) \cap L^1_{\text{loc}}(\Omega)$ and let $u \in H^1_0(\Omega)$ be such that a.e. on Ω

$$\text{Re } T \cdot \bar{u} \geq f$$

for some real valued function $f \in L^1(\Omega)$. Then $\text{Re } T \cdot \bar{u} \in L^1(\Omega)$ and

$$\text{Re} \langle T, u \rangle = \int \text{Re } T \cdot \bar{u}$$

where $\langle T, u \rangle$ denotes the Hermitian scalar product in the duality between $H^{-1}(\Omega)$ and $H^1_0(\Omega)$.

The proof of Theorem 2.1 is divided into 4 steps.

Step 1: $A + \lambda$ is onto for $\lambda > \lambda_1$. Set $q_n^+ = \min(q^+, n)$; by a Theorem of Lax-Milgram there exists a unique function $u_n \in H^1_0(\Omega)$ which satisfies

$$(4) \quad -\Delta u_n + (q_n^+ - q^-)u_n + \lambda u_n = f.$$

⁽¹⁾ The use of this sort of lemma in this context was suggested by M. Crandall.

(Note that by Lemma 2.1 the sesquilinear form $\int q^- u \bar{v}$ is continuous on $H_0^1(\Omega)$).

Multiplying (4) by \bar{u}_n we find a constant C independent of n such that

$$(5) \quad \|u_n\|_{H^1} \leq C,$$

$$(6) \quad \int q_n^+ |u_n|^2 \leq C.$$

Choose a subsequence denoted again by u_n such that $u_n \rightarrow u$ weakly in $H_0^1(\Omega)$ and $u_n \rightarrow u$ a.e. on Ω . It follows from Fatou's Lemma and (6) that $q^+ |u|^2 \in L^1(\Omega)$. We deduce that $qu \in L_{loc}^1(\Omega)$; indeed

$$q^+ |u| \leq \frac{1}{2} q^+ (|u|^2 + 1) \in L_{loc}^1(\Omega),$$

$$q^- |u| \leq \frac{1}{2} q^- (|u|^2 + 1) \in L_{loc}^1(\Omega).$$

We pass now to the limit in (4) and prove that $-\Delta u + qu + \lambda u = f$ in $D'(\Omega)$. It suffices to show that

$$(q_n^+ - q^-) u_n + qu \text{ in } L_{loc}^1(\Omega).$$

For this purpose we adapt a device due to W. Strauss [9] and extensively used in the study of strongly nonlinear equations. In view of Vitali's convergence theorem, it suffices to verify that given $\omega \subset\subset \Omega$, then $\forall \epsilon > 0, \exists \delta > 0$ such that $E \subset \omega$ and $|E| < \delta$ imply $\int_E |q_n^+ - q^-| |u_n| < \epsilon$ for all n . But for every $R > 0$ we have

$$q_n^+ |u_n| \leq \frac{1}{2} q_n^+ (R + \frac{1}{R} |u_n|^2)$$

and thus, by (6),

$$\int_E q_n^+ |u_n| \leq \frac{1}{2} R \int_E q^+ + \frac{1}{2R} C.$$

We fix R large enough so that $\frac{C}{R} < \epsilon$ and then $\delta > 0$ so small that $R \int_E q^+ < \epsilon$. We proceed similarly with $q^- |u_n|$.

Step 2: $A + \lambda_1$ is accretive. Let $u \in D(A)$ and set $T = qu$. Since $T \in H^{-1}(\Omega) \cap L_{loc}^1(\Omega)$ and

$$\operatorname{Re} T \bar{u} = q |u|^2 \geq -q^- |u|^2 \in L^1(\Omega)$$

it follows from Lemma 2.2 that $q |u|^2 \in L^1$ and

$$\operatorname{Re} \langle T, u \rangle = \int q |u|^2.$$

But $qu = Au + \lambda u$ and so

$$\operatorname{Re}(Au, u) - \int |\operatorname{grad} u|^2 = \int q|u|^2.$$

Since $Au \in L^2(\Omega)$ we have in fact

$$\operatorname{Re}(Au, u) = \int |\operatorname{grad} u|^2 + \int q|u|^2 \geq -\lambda_1 \int |u|^2$$

by Lemma 2.1.

Step 3: $u \in D(A)$ implies $q|u|^2 \in L^1(\Omega)$ and (2) holds. We have just seen in Step 2 that $u \in D(A)$ implies $q|u|^2 \in L^1(\Omega)$. Now let $u, v \in D(A)$ and set $T = qu$. We have $T \in H^{-1}(\Omega) \cap L^1_{\text{loc}}(\Omega)$ and

$$\operatorname{Re} T \cdot \bar{v} = \operatorname{Re} qu\bar{v} \geq -\frac{1}{2} |q||u|^2 - \frac{1}{2} |q||v|^2 \in L^1(\Omega)$$

and therefore

$$\operatorname{Re}(T, v) = \int \operatorname{Re} qu\bar{v}.$$

Thus

$$\operatorname{Re}(Au, v) - \operatorname{Re} \int \operatorname{grad} u \operatorname{grad} \bar{v} = \operatorname{Re} \int qu\bar{v}.$$

Changing u into iu we find

$$(Au, v) = \int \operatorname{grad} u \operatorname{grad} \bar{v} + \int qu\bar{v}.$$

Step 4: A is self-adjoint. Indeed $A + \lambda_1$ is m -accretive and symmetric. Therefore $A + \lambda_1$ is self-adjoint and so is A .

Proof of Theorem 2.2: Clearly $A \subset A_1$. Let $u \in D(A_1)$ and set $f = A_1 u + \lambda u$ with some $\lambda > \lambda_1$. Let $u^* \in D(A)$ be the unique solution of

$$Au^* + \lambda u^* = f.$$

We have

$$A_1(u - u^*) + \lambda(u - u^*) = 0.$$

Since $(u - u^*) \in L^1_{\text{loc}}(\mathbb{R}^m)$ and $\Delta(u - u^*) \in L^1_{\text{loc}}(\mathbb{R}^m)$ we may apply Lemma A in [3] to conclude that

$$\Delta|u - u^*| \geq \operatorname{Re}[\Delta(u - u^*) \operatorname{sign}(\bar{u} - \bar{u}^*)] \text{ in } D'(\mathbb{R}^m),$$

and thus in $D'(\mathbb{R}^m)$ we find,

$$\Delta|u - u^*| \geq \operatorname{Re}[(q + \lambda)|u - u^*|] \geq (-q^- + \lambda)|u - u^*|.$$

Using the next lemma we conclude that $u = u^*$ (and hence $D(A_1) = D(A)$).

Lemma 2.3: Assume (1) and (3). Let $v \in L^2(\mathbb{R}^m)$ be a real valued function with $q^- v \in L^1_{loc}(\mathbb{R}^m)$ satisfying

$$-\Delta v - q^- v + \lambda v \leq 0 \text{ in } D'(\mathbb{R}^m)$$

with some $\lambda > \lambda_1$. Then $v \leq 0$ a.e. on \mathbb{R}^m .

The proof of Lemma 2.3 relies on the following crucial result. Since we shall need it in Section 3 for a general domain $\Omega \subset \mathbb{R}^m$ we work now again in Ω .

Theorem 2.3: Assume (1). Let $g \in L^2(\Omega) \cap L^\infty(\Omega)$ and let $\psi \in H^1_0(\Omega)$ be the unique solution of

$$(7) \quad -\Delta \psi - q^- \psi + \lambda \psi = g \text{ in } \Omega \quad (\lambda > \lambda_1).$$

Then

a) $g \geq 0$ a.e. on Ω implies $\psi \geq 0$ a.e. on Ω ;

b) $\psi \in \bigcap_{2 \leq p < \infty} L^p(\Omega)$.

Proof of Theorem 2.3: a) Multiplying (7) by $-\psi^-$ we find

$$\int |\text{grad } \psi^-|^2 - \int q^- |\psi^-|^2 + \lambda \int |\psi^-|^2 \leq 0$$

and thus $\psi^- = 0$.

b) We have to consider only the case $m \geq 3$ (when $m \leq 2$, $\psi \in H^1_0(\Omega)$ implies

$\psi \in \bigcap_{2 \leq p < \infty} L^p(\Omega)$).

We can always assume that $g \geq 0$ a.e. on Ω so that $\psi \geq 0$ a.e. on Ω . We truncate q^- by $q_k^- = \min(q^-, k)$ and define ψ_k to be the unique solution of

$$\begin{cases} \psi_k \in H^1_0(\Omega) \\ -\Delta \psi_k - q_k^- \psi_k + \lambda \psi_k = g \text{ in } \Omega. \end{cases}$$

It is clear that $\psi_k \rightarrow \psi$ weakly in $H^1_0(\Omega)$ as $k \rightarrow \infty$. We shall prove that for every $p \in [2, \infty)$, $\psi_k \in L^p(\Omega)$ and

$$(8) \quad \|\psi_k\|_{L^p} \leq C_p (\|g\|_{L^2} + \|g\|_{L^\infty}),$$

where C_p is independent of k , but it depends on q^- through the use of Lemma 2.1.

For simplicity we drop now the subscript k on ψ_k and write

$$(9) \quad -\Delta \psi - q_k^- \psi + \lambda \psi = g.$$

Set $\psi_n = \min(\psi, n)$ and let $2 \leq p < \infty$; since $(\psi_n)^{p-1} \in H_0^1(\Omega)$ we can multiply (9) by $(\psi_n)^{p-1}$ and we get

$$(p-1) \int (\psi_n)^{p-2} |\text{grad } \psi_n|^2 \leq \int g(\psi_n)^{p-1} + \int q_k^-(\psi_n)^p + \int_{[\psi > n]} kn^{p-1} \psi,$$

that is

$$\begin{aligned} \frac{4(p-1)}{p^2} \int |\text{grad } \psi_n^{p/2}|^2 &\leq \|g\|_{L^p} \|\psi_n\|_{L^p}^{p-1} + \int q_k^-(\psi_n)^p + kn^{p-1} \int_{[\psi > n]} \psi \\ &\leq \|g\|_{L^p} \|\psi_n\|_{L^p}^{p-1} + \epsilon \|\text{grad } \psi_n^{p/2}\|_{L^2}^2 + \lambda_\epsilon \|\psi_n\|_{L^p}^p + k \int_{[\psi > n]} \psi^p \end{aligned}$$

by Lemma 2.1 (here $\int_{[\psi > n]} \psi^p$ is possibly infinite). Choosing $\epsilon > 0$ small enough (for example $\epsilon = \frac{2(p-1)}{p^2}$) we see that

$$\int |\text{grad } \psi_n^{p/2}|^2 \leq C_p [\|g\|_{L^p}^p + \|\psi\|_{L^p}^p + k \int_{[\psi > n]} \psi^p]$$

where C_p is independent of k and n . Using Sobolev's inequality we find

$$(10) \quad \|\psi\|_{L^{p^2/2}}^p \leq C_p \left[\|g\|_{L^p}^p + \|\psi\|_{L^p}^p + k \int_{[\psi > n]} \psi^p \right].$$

Assuming now that $\psi \in L^p(\Omega)$ and passing to the limit in (10) as $n \rightarrow \infty$ we obtain that $\psi \in L^{p^2/2}(\Omega)$ and

$$\|\psi\|_{L^{p^2/2}} \leq C_p [\|g\|_{L^p} + \|\psi\|_{L^p}].$$

Iterating this process from $p = 2$ we obtain finally for every $p \in [2, \infty)$

$$\|\psi\|_{L^p} \leq C_p [\|g\|_{L^2} + \|g\|_{L^\infty}].$$

More precisely we have proved (8). The conclusion of Theorem 2.3 follows since $\psi_k \rightarrow \psi$ weakly in $H_0^1(\Omega)$ as $k \rightarrow \infty$.

Proof of Lemma 2.3: By assumption $q^-v \in L_{loc}^1(\mathbb{R}^m)$ and

$$\int v(-\Delta\varphi - q^-v + \lambda\varphi) \leq 0 \quad \forall \varphi \in D_+(\mathbb{R}^m).$$

An easy density argument (smoothing by convolution) shows that

$$(11) \quad \int v(-\Delta\varphi - q^-v + \lambda\varphi) \leq 0 \quad \forall \varphi \in H^2(\mathbb{R}^m) \cap L^\infty(\mathbb{R}^m), \text{ supp } \varphi \text{ compact, } \varphi \geq 0 \text{ a.e.}$$

Fix $g \in D_+(\mathbb{R}^m)$ and let $\psi_k \in H^1(\mathbb{R}^m)$ be the unique solution of

$$(12) \quad -\Delta \psi_k - q_k^- \psi_k + \lambda \psi_k = g \text{ in } \mathbb{R}^m.$$

We know by Theorem 2.3 that $\psi_k \geq 0$ a.e.

$$\psi_k \in \bigcap_{2 \leq p < \infty} L^p(\mathbb{R}^m) \text{ with } \|\psi_k\|_{L^p} \leq C_p,$$

and also $\|\text{grad } \psi_k\|_{L^2} \leq C$. In addition we derive from (12) that

$$\psi_k \in H^2(\mathbb{R}^m) \cap L_{loc}^\infty(\mathbb{R}^m).$$

Fix $\zeta \in D_+(\mathbb{R}^m)$ satisfying $\zeta(x) = 1$ for $|x| \leq 1$ and set $\zeta_n(x) = \zeta(\frac{x}{n})$. In (11)

we choose $\varphi = \psi_k \zeta_n$. Note that by (12)

$$-\Delta \varphi - q^- \varphi + \lambda \varphi = \zeta_n g - (\Delta \zeta_n) \psi_k - 2 \text{grad } \zeta_n \text{ grad } \psi_k - \zeta_n \psi_k (q^- - q_k^-),$$

and therefore

$$\int v \zeta_n g \leq \frac{C}{2} + \frac{C}{n} + \int v \zeta_n \psi_k (q^- - q_k^-).$$

First we fix n and let $k \rightarrow \infty$. We distinguish two cases:

a) $m \geq 5$,

b) $m < 5$.

a) When $m \geq 5$ we have $q^- - q_k^- \rightarrow 0$ in $L_{loc}^{m/2}(\mathbb{R}^m)$. Let $p \in [2, \infty)$ be such that

$\frac{1}{2} + \frac{2}{m} + \frac{1}{p} = 1$; we have

$$|\int v \zeta_n \psi_k (q^- - q_k^-)| \leq \|v\|_{L^2} \|\psi_k\|_{L^p} \|\zeta_n (q^- - q_k^-)\|_{L^{m/2}} \rightarrow 0.$$

Consequently

$$\int v \zeta_n g \leq \frac{C}{2} + \frac{C}{n}.$$

b) When $m < 5$ we use the assumption (3) (or (1)): $q^- \in L_{loc}^{m/2+\epsilon}(\mathbb{R}^m)$ with some $\epsilon > 0$.

It follows from (12) that ψ_k remains bounded in $W_{loc}^{2,q}(\mathbb{R}^m)$ for some $q > \frac{m}{2}$ (when

$m \geq 2$) as $k \rightarrow \infty$. We conclude that ψ_k remains bounded in $L_{loc}^\infty(\mathbb{R}^m)$ as $k \rightarrow \infty$

(in case $m = 1$, ψ_k is bounded in $L^\infty(\mathbb{R})$ since it is bounded in $H^1(\mathbb{R})$). Therefore

$$\int v \zeta_n \psi_k (q^- - q_k^-) \rightarrow 0 \text{ as } k \rightarrow \infty$$

since $\|\zeta_n v (q^- - q_k^-)\|_{L^1} \rightarrow 0$ by the dominated convergence theorem (recall that

$q^- v \in L^1_{loc}(\mathbb{R}^m)$. In both cases we find

$$\int v \zeta_n g \leq \frac{C}{2} + \frac{C}{n} v_n.$$

As $n \rightarrow \infty$ we see that

$$\int v g \leq 0 \quad \forall g \in D_+(\mathbb{R}^m)$$

and therefore $v \leq 0$ a.e. on \mathbb{R}^m .

Remark 2.2: The conclusion of Lemma 2.3 fails in \mathbb{R}^3 and in \mathbb{R}^4 if we do not assume

(3). Ancona (personal communication) has constructed in \mathbb{R}^3 and in \mathbb{R}^4 functions

$q^- \in L^{m/2}(\mathbb{R}^m)$ and $u \in L^{m/m-2}(\mathbb{R}^m) \cap L^2(\mathbb{R}^m)$ such that

$$-\Delta u - q^- u + u = 0 \text{ in } D'$$

with $\|q^-\|_{L^{m/2}}$ as small as we please and $u \neq 0$.

3. Complex potentials

Let Ω be an (arbitrary) open subset of \mathbb{R}^m . Assume $q(x)$ and $q'(x)$ are real valued functions such that $q, q' \in L^1_{loc}(\Omega)$ and set

$$V(x) = q(x) + iq'(x).$$

We assume

$$(13) \quad \text{either } q' \in L^{1+\varepsilon}_{loc}(\Omega) \text{ or } q^- \in L^{(m/2)+\varepsilon}_{loc}(\Omega) \text{ when } m \geq 2,$$

for some arbitrarily small $\varepsilon > 0$. Define

$$A = -\Delta + V(x)$$

with

$$D(A) = \{u \in H^1_0(\Omega); Vu \in L^1_{loc}(\Omega) \text{ and } -\Delta u + Vu \in L^2(\Omega)\}.$$

The main results are the following

Theorem 3.1: Assume (1) and (13). Then A is closable in $L^2(\Omega)$ and $\bar{A} + \lambda_1$ is m -accretive. In addition $u \in D(\bar{A})$ implies that $u \in H^1_0(\Omega)$, $q|u|^2 \in L^1(\Omega)$ and

$$(14) \quad \operatorname{Re}(\bar{A}u, u) = \int |\operatorname{grad} u|^2 + \int q|u|^2.$$

Remark 3.1: In case we assume

$$(15) \quad |q'(x)| \leq Mq^+(x) + h(x) \text{ for a.e. } x \in \Omega$$

with $h \in L^{2m/(m+2)}_{loc}(\Omega)$ and $m \geq 3$ then A is closed in $L^2(\Omega)$. (Note that (15) corresponds essentially with the assumption made in [5]). Indeed let $u_n \in D(A)$ be such that

$u_n \rightarrow u$ in $L^2(\Omega)$ and $Au_n \rightarrow f$ in $L^2(\Omega)$. It follows from Lemma 2.1 and (14) that $u_n \rightarrow u$ in $H^1_0(\Omega)$ and $\sqrt{q^+}u_n \rightarrow \sqrt{q^+}u$ in $L^2(\Omega)$. From (15) we deduce easily that $Vu \in L^1_{loc}(\Omega)$ and that $-\Delta u + Vu = f$ in $D'(\Omega)$. Therefore $u \in D(A)$ and $Au = f$.

When $\Omega = \mathbb{R}^m$ we consider also the operator A_1 defined in $L^2(\mathbb{R}^m)$ by

$$A_1 = -\Delta + V(x)$$

with

$$D(A_1) = \{u \in L^2(\mathbb{R}^m); Vu \in L^1_{loc}(\mathbb{R}^m) \text{ and } -\Delta u + Vu \in L^2(\mathbb{R}^m)\}.$$

Theorem 3.2: Assume (1), (3) and (13). Then A_1 is closable and $\bar{A}_1 = \bar{A}$.

In the proof of Theorem 3.1 we shall use the following

Lemma 3.1: Let $v \in H_0^1(\Omega)$ be a real valued function. Assume (1) and

$$-\Delta v - q^- v + \lambda v \leq 0 \text{ in } D'(\Omega)$$

with $\lambda > \lambda_1$. Then $v \leq 0$ a.e. on Ω .

Proof of Lemma 3.1: We have, for every $\varphi \in D_+(\Omega)$

$$\int \text{grad} v \text{ grad} \varphi - \int q^- v \varphi + \lambda \int v \varphi \leq 0.$$

Now we use the fact (pointed out by G. Stampacchia) that $D_+(\Omega)$ is dense in

$\{u \in H_0^1(\Omega); u \geq 0 \text{ a.e. on } \Omega\}$ for the H^1 norm⁽¹⁾ to derive that

$$\int \text{grad} v \text{ grad} \varphi - \int q^- v \varphi + \lambda \int v \varphi \leq 0 \quad \forall \varphi \in H_0^1(\Omega), \varphi \geq 0.$$

Choosing $\varphi = v^+$ we obtain

$$\int |\text{grad} v^+|^2 - \int q^- |v^+|^2 + \lambda \int |v^+|^2 \leq 0$$

and therefore $v^+ = 0$.

The proof of Theorem 3.1 is divided into five steps.

Step 1: $R(A + \lambda) \supset L^2(\Omega) \cap L^\infty(\Omega)$ for $\lambda > \lambda_1$.

Indeed let $f \in L^2(\Omega) \cap L^\infty(\Omega)$ and let $u_n \in H_0^1(\Omega)$ be the unique solution of

$$(16) \quad -\Delta u_n + V_n u_n + \lambda u_n = f$$

where $V_n = q_n^+ - q^- + i q_n'$ and

$$q_n' = \begin{cases} n & \text{if } q' > n \\ q' & \text{if } |q'| \leq n \\ -n & \text{if } q' \leq -n. \end{cases}$$

The existence of u_n follows from a theorem of Lax-Milgram. Multiplying (16) by $\overline{u_n}$ we find

$$(17) \quad \|u_n\|_{H^1} \leq C$$

$$(18) \quad \int q_n^+ |u_n|^2 \leq C.$$

(1) Indeed let $u \in H_0^1(\Omega)$ with $u \geq 0$ a.e. on Ω ; let $u_n \in D(\Omega)$ be such that $u_n \rightarrow u$ in $H^1(\Omega)$. We claim that $|u_n| \rightarrow |u| = u$ in $H^1(\Omega)$ because $\| |u_n| \|_{H^1} = \| u_n \|_{H^1}$ and $|u_n| \rightarrow |u|$ weakly in $H^1(\Omega)$. On the other hand $|u_n|$ can be smoothed by convolution and for fixed n , $\rho_\epsilon * |u_n| \rightarrow |u_n|$ in $H^1(\Omega)$ as $\epsilon \rightarrow 0$.

On the other hand we have

$$\Delta |u_n| \geq \operatorname{Re}[\Delta u_n \operatorname{sign} \overline{u_n}] \quad \text{in } D'(\Omega)$$

which leads to

$$-\Delta |u_n| - q^- |u_n| + \lambda |u_n| \leq |f| \quad \text{in } D'(\Omega).$$

Let $\psi \in H_0^1(\Omega)$ be the solution of

$$(19) \quad -\Delta \psi - q^- \psi + \lambda \psi = |f|.$$

It follows from Lemma 3.1 that

$$(20) \quad |u_n| \leq \psi \quad \text{a.e. on } \Omega.$$

By Theorem 2.3 we know that $\psi \in L^p(\Omega)$ for every $p \in [2, \infty)$. We extract a subsequence, denoted again by u_n such that $u_n \rightharpoonup u$ weakly in $H_0^1(\Omega)$, $u_n \rightarrow u$ a.e. on Ω . We see as in the proof of Theorem 2.1 (Step 1) that $(q_n^+ - q^-)u_n \rightarrow qu$ in $L_{loc}^1(\Omega)$. Therefore we have only to verify that $q_n' u_n \rightarrow q'u$ in $L_{loc}^1(\Omega)$. We distinguish two cases:

a) $q' \in L_{loc}^{1+\varepsilon}(\Omega)$,

b) $q^- \in L_{loc}^{(m/2)+\varepsilon}(\Omega)$.

Case a) From (20) we deduce that $u_n \rightarrow u$ in every L^p space, $2 \leq p < \infty$ and so

$$q_n' u_n \rightarrow q'u \quad \text{in } L_{loc}^1(\Omega).$$

Case b) Since $q^- \psi \in L_{loc}^q(\Omega)$ for some $q > \frac{m}{2}$, it follows from (19) that $\psi \in L_{loc}^\infty(\Omega)$.

We deduce from the dominated convergence theorem that $q_n' u_n \rightarrow q'u$ in $L_{loc}^1(\Omega)$.

Step 2: $A + \lambda_1$ is accretive. Let $u \in D(A)$ and set $T = Vu$. We have

$$T \in H^{-1}(\Omega) \cap L_{loc}^1(\Omega) \quad \text{and}$$

$$\operatorname{Re} T \cdot \bar{u} = q|u|^2 \geq -q^- |u|^2 \in L^1(\Omega).$$

It follows from Lemma 2.2 that $q|u|^2 \in L^1(\Omega)$ and

$$\int q|u|^2 = \operatorname{Re}(T, u) = \operatorname{Re}(Au + \Delta u, u).$$

Therefore

$$(21) \quad \operatorname{Re}(Au, u) = \int |\operatorname{grad} u|^2 + \int q|u|^2 \geq -\lambda_1 \int |u|^2.$$

Step 3: $D(A)$ is dense in $L^2(\Omega)$. Given $f \in L^2(\Omega) \cap L^\infty(\Omega)$ we solve for large n the equation

$$(22) \quad u_n + \frac{1}{n} Au_n = f.$$

We shall prove that $u_n \rightarrow f$ in $L^2(\Omega)$ as $n \rightarrow \infty$ — and as a consequence $D(A)$ is dense in $L^2(\Omega)$. By (21) we have

$$\int |u_n|^2 + \frac{1}{n} \int |\text{grad} u_n|^2 + \frac{1}{n} \int q |u_n|^2 = \text{Re}(f, u_n).$$

In particular we deduce that

$$(23) \quad \limsup_{n \rightarrow \infty} \|u_n\|_{L^2} \leq \|f\|_{L^2}$$

$$(24) \quad \frac{1}{n} \int q^+ |u_n|^2 \leq c$$

$$(25) \quad \frac{1}{n} \int |\text{grad} u_n|^2 \leq c.$$

Next we have (as in the proof of Step 1)

$$|u_n| - \frac{1}{n} \Delta |u_n| - \frac{1}{n} q^- |u_n| \leq |f| \quad \text{in } D'(\Omega).$$

On the other hand let $\psi \in H_0^1(\Omega)$ be the solution of

$$-\Delta \psi - q^- \psi + \lambda \psi = |f|$$

for some fixed $\lambda > \lambda_1$. Since $|u_n| \geq \lambda \left| \frac{u_n}{n} \right|$ for $n \geq \lambda$, we deduce from Lemma 3.1 that $\left| \frac{u_n}{n} \right| \leq \psi$ a.e. Choose a subsequence, denoted again by u_n such that $u_n \rightarrow u$ weakly in $L^2(\Omega)$, $\frac{1}{n} u_n \rightarrow 0$ a.e. (this is possible since $\frac{1}{n} u_n \rightarrow 0$ in $L^2(\Omega)$). For every $\varphi \in D(\Omega)$ we have

$$(26) \quad \int u_n \bar{\varphi} - \frac{1}{n} \int u_n \Delta \bar{\varphi} + \frac{1}{n} \int v u_n \bar{\varphi} = \int f \bar{\varphi}.$$

We claim that $\frac{1}{n} \int v u_n \bar{\varphi} \rightarrow 0$ as $n \rightarrow \infty$. Indeed by (24) and (25) we have

$$\frac{1}{n} \left| \int q^+ u_n \bar{\varphi} \right| \leq \frac{C}{\sqrt{n}} \quad \text{and} \quad \frac{1}{n} \left| \int q^- u_n \bar{\varphi} \right| \leq \frac{C}{\sqrt{n}}.$$

Thus we have only to verify that $\frac{1}{n} \int q' u_n \bar{\varphi} \rightarrow 0$. We distinguish two cases:

- a) if $q^- \in L_{\text{loc}}^{(m/2)+\epsilon}(\Omega)$, we have $\psi \in L_{\text{loc}}^\infty(\Omega)$ and we deduce from the dominated convergence theorem that $\frac{1}{n} \int q' u_n \bar{\varphi} \rightarrow 0$;
- b) if $q' \in L_{\text{loc}}^{1+\epsilon}(\Omega)$ we use the fact that $\left| \frac{u_n}{n} \right| \leq \psi \in L^p(\Omega)$ for every $2 \leq p < \infty$ to deduce that $\frac{u_n}{n} \rightarrow 0$ in $L^p(\Omega)$ and so $\frac{1}{n} \int q' u_n \bar{\varphi} \rightarrow 0$.

In all the cases, we derive from (26) that

$$\int u \bar{\psi} = \int f \bar{\psi} \quad \forall \psi \in D$$

and consequently $u = f$. We conclude using (23) that $u_n \rightarrow f$ in $L^2(\Omega)$.

Step 4: A is closable and $\bar{A} + \lambda_1$ is m -accretive. This is a standard fact, see e.g. Theorem 3.4 in [2].

Step 5: $u \in D(\bar{A})$ implies that $u \in H_0^1(\Omega)$, $q|u|^2 \in L^1(\Omega)$ and (14).

We already know (Step 2) that $v \in D(A)$ implies $q|v|^2 \in L^1(\Omega)$ and

$$(27) \quad \operatorname{Re}(Av, v) = \int |\operatorname{grad} v|^2 + \int q|v|^2.$$

Now let $u \in D(\bar{A})$ and let $u_n \in D(A)$ be such that $u_n \rightarrow u$, $Au_n \rightarrow \bar{A}u$. It follows from

(27) applied to $v = u_n - u_m$ that $u_n \rightarrow u$ in $H_0^1(\Omega)$ and $\int q^+ |u_n - u|^2 \rightarrow 0$ (since u_n is a Cauchy sequence in $H_0^1(\Omega)$ and in $L^2(\Omega)$ with weight q^+). In particular $q|u|^2 \in L^1(\Omega)$ and (14) holds.

Proof of Theorem 3.2: Clearly $A \subset A_1$. Now let $u \in D(A_1)$ and let $\lambda > \lambda_1$. Set $f = A_1 u + \lambda u$, and let u^* be the unique solution of

$$\bar{A}u^* + \lambda u^* = f.$$

Thus, there exists a sequence $u_n^* \rightarrow u^*$ in $L^2(\mathbb{R}^m)$ with $u_n^* \in D(A)$ and

$$Au_n^* + \lambda u_n^* = f_n + f \quad \text{in } L^2(\mathbb{R}^m).$$

In particular we have

$$A_1(u_n^* - u) + \lambda(u_n^* - u) = f_n - f$$

and therefore

$$-\Delta |u_n^* - u| - q^- |u_n^* - u| + \lambda |u_n^* - u| \leq |f_n - f| \quad \text{in } D'(\mathbb{R}^m).$$

We deduce from Lemma 2.3 that $|u_n^* - u| \leq \psi_n$ a.e. on \mathbb{R}^m where $\psi_n \in H^1(\mathbb{R}^m)$ is the solution of

$$-\Delta \psi_n - q^- \psi_n + \lambda \psi_n = |f_n - f|.$$

Hence $\|\psi_n\|_{H^1} \rightarrow 0$ and in particular $u_n^* - u \rightarrow 0$ in $L^2(\mathbb{R}^m)$. It follows that $u^* = u$, that is $A_1 \subset \bar{A}$. We have $A \subset A_1 \subset \bar{A}$ and therefore A_1 is closable with $\bar{A}_1 = \bar{A}$.

REFERENCES

- [1] H. Brezis and F. Browder, Sur une propriété des espaces de Sobolev, C. R. Acad. Sc. Paris (1978).
- [2] T. Kato, Perturbation theory for linear operators, 2nd edition, Springer (1976).
- [3] T. Kato, Schrödinger operators with singular potentials, Israel J. Math. 13 (1972), p. 135-148.
- [4] T. Kato, A second look at the essential selfadjointness of the Schrödinger operator, Physical Reality and Mathematical Description, D. Reidel Publishing Co. (1976), p. 193-201.
- [5] T. Kato, On some Schrödinger operators with a singular complex potential, Ann. Sc. Norm. Sup. Pisa Ser. IV, 5 (1978), p. 105-114.
- [6] E. Nelson, Feynman integrals and the Schrödinger equation, J. Math. Phys. 5 (1964), p. 332-343.
- [7] C. Simader, Bemerkungen über Schrödinger-Operatoren mit stark singularen Potentialen, Math. Z. 138 (1974), p. 53-70.
- [8] B. Simon, Essential self-adjointness of Schrödinger operators with positive potentials, Math. Ann. 201 (1973), p. 211-220.
- [9] W. Strauss, On weak solutions of semilinear hyperbolic equations, Ann. Acad. Bras. Cienc. 42 (1970), p. 645-651.

REPORT DOCUMENTATION PAGE		READ INSTRUCTIONS BEFORE COMPLETING FORM
1. REPORT NUMBER 1875	2. GOVT ACCESSION NO.	3. RECIPIENT'S CATALOG NUMBER
4. TITLE (and Subtitle) REMARKS ON THE SCHRÖDINGER OPERATOR WITH SINGULAR COMPLEX POTENTIALS		5. TYPE OF REPORT & PERIOD COVERED Summary Report - no specific reporting period
7. AUTHOR(s) Haïm Brezis and Tosio Kato		6. PERFORMING ORG. REPORT NUMBER
9. PERFORMING ORGANIZATION NAME AND ADDRESS Mathematics Research Center, University of Wisconsin 610 Walnut Street Madison, Wisconsin 53706		8. CONTRACT OR GRANT NUMBER(s) DAAG29-75-C-0024 MCS76-04655
11. CONTROLLING OFFICE NAME AND ADDRESS See Item 18 below.		10. PROGRAM ELEMENT, PROJECT, TASK AREA & WORK UNIT NUMBERS Work Unit Number 1 - Applied Analysis
14. MONITORING AGENCY NAME & ADDRESS (if different from Controlling Office)		12. REPORT DATE August 1978
		13. NUMBER OF PAGES 17
		15. SECURITY CLASS. (of this report) UNCLASSIFIED
		15a. DECLASSIFICATION/DOWNGRADING SCHEDULE
16. DISTRIBUTION STATEMENT (of this Report) Approved for public release; distribution unlimited.		
17. DISTRIBUTION STATEMENT (of the abstract entered in Block 20, if different from Report)		
18. SUPPLEMENTARY NOTES U. S. Army Research Office P. O. Box 12211 Research Triangle Park North Carolina 27709 National Science Foundation Washington, D. C. 20550		
19. KEY WORDS (Continue on reverse side if necessary and identify by block number) Schrödinger operator Complex potentials m-accretive operator		
20. ABSTRACT (Continue on reverse side if necessary and identify by block number) Let $A = -\Delta + V(x)$ be a Schrödinger operator on an (arbitrary) open set $\Omega \subset \mathbb{R}^m$, where $V \in L^1_{loc}(\Omega)$ is a complex valued function. We consider the "maximal" realization of A in $L^2(\Omega)$ under Dirichlet boundary condition, that is		

20. ABSTRACT - cont'd.

$$D(A) = \{u \in H_0^1(\Omega); Vu \in L_{loc}^1(\Omega) \text{ and } -\Delta u + Vu \in L^2(\Omega)\}.$$

When $\Omega = \mathbb{R}^m$ we also consider the operator

$$A_1 = -\Delta + V$$

with domain

$$D(A_1) = \{u \in L^2(\Omega); Vu \in L_{loc}^1(\Omega) \text{ and } -\Delta u + Vu \in L^2(\Omega)\}.$$

A special case of our main results is:

Theorem: Let $m \geq 3$; assume that the function $\max\{-\operatorname{Re} V, 0\}$ belongs to $L^\infty(\Omega) + L^{m/2}(\Omega)$ and also to $L_{loc}^{(m/2)+\epsilon}(\Omega)$ for some $\epsilon > 0$. Then A (resp. A_1) is closable and $\bar{A} + \lambda$ (resp. $\bar{A}_1 + \lambda$) is m -accretive for some real constant λ .